

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



09/597, 102

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07C 211/00</b>		A2	(11) International Publication Number: <b>WO 99/01416</b>
			(43) International Publication Date: 14 January 1999 (14.01.99)
(21) International Application Number: PCT/EP98/04026			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO <sup>7</sup> patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 24 June 1998 (24.06.98)			
(30) Priority Data: 08/885,944 30 June 1997 (30.06.97) US			
(71) Applicant: MERZ + CO. GMBH & CO. [DE/DE]; Eckenheimer Landstrasse 100-104, D-60318 Frankfurt am Main (DE).			
(72) Inventors: GOLD, Markus; Kranichstrasse 9, D-6085 Nauheim (DE). DANYSZ, Wojciech; Vor den Garten 16, D-61130 Nidderau (DE). PARSONS, Christopher, Graham, Raphael; Alt Praunheim 99, D-60488 Praunheim (DE). KALVINSH, Ivars; Miera 17-8, LV-2169 Salaspils (LV). KAUSS, Valerjans; A. Deglava 108/2-36, LV-1082 Riga (LV). JIRGENSONS, Aigars; Basteja Bulv. 10-48, LV-1050 Riga (LV).			
(74) Agent: MARSDEN, John, Christopher; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).			Published Without international search report and to be republished upon receipt of that report.
(54) Title: 1-AMINO-ALKYLCYCLOHEXANE NMDA RECEPTOR ANTAGONISTS			
(57) Abstract <p>Certain 1-amino-alkylcyclohexanes are systemically-active uncompetitive NMDA receptor antagonists having rapid blocking/unblocking kinetics and strong voltage-dependency and are therefore useful in the alleviation of conditions resulting from disturbances of glutamatergic transmission giving them a wide range of utility in the treatment of CNS disorders involving the same, as well as in non-NMDA indications, due to their immunomodulatory, antimalarial, anti-Borna virus, and anti-Hepatitis C activities and utilities. Pharmaceutical compositions thereof and a method of treating conditions which are alleviated by the employment of an NMDA receptor antagonist, as well as the aforementioned non-NMDA indications, and a method for the preparation of the active 1-amino-alkylcyclohexane compounds involved.</p>			

102  
inherent  
claim 12-23

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

1-AMINO-ALKYLCYCLOHEXANE NMDA RECEPTOR ANTAGONISTSBACKGROUND OF THE INVENTION

## 1. Field of the Invention

1-Amino-alkylcyclohexane compounds which are systemically-active as NMDA receptor antagonists, pharmaceutical compositions comprising the same, method of preparation thereof, and method of treating CNS disorders which involve disturbances of glutamatergic transmission therewith.

## 2. Prior Art

Antagonism of glutamate receptors of the N-methyl-D-aspartate (NMDA) type has a potentially wide range of therapeutic applications [19]. Functional inhibition of NMDA receptors can be achieved through actions at different recognition sites such as the primary transmitter site, strychnine-insensitive glycine site (glycine<sub>B</sub>), polyamine site, and phencyclidine site located inside the cation channel. The NMDA receptor channel blockers act in an uncompetitive "use-dependent" manner, meaning that they usually only block the channel in the open state. This use-dependence has been interpreted by many to mean that stronger activation of the receptor should lead to a greater degree of antagonism. Such a mode of action has further been taken to imply that this class of antagonist may be particularly useful when overactivation of NMDA receptors can be expected, such as in epilepsy, ischaemia, and trauma. However, initial clinical experience with the selective, high affinity, strongly use-dependent uncompetitive NMDA receptor antagonist (+)-5-methyl-10,11-dihydro-

5H-dibenzocyclohepten-5,10-imine maleate ((+)-MK-801) has been disappointing. Namely, therapeutic efficacy in epilepsy was poor while some psychotropic side effects were apparent at therapeutic doses. These observations, together with the fact that phencyclidine abusers experience similar psychotropic symptoms, has led to the conclusion that uncompetitive antagonism of NMDA receptors may not be a promising therapeutic approach.

However, the use of more elaborate electrophysiological methods indicates that there is no equality between different uncompetitive antagonists since factors such as the speed of receptor blockade (on-off kinetics) and the voltage-dependence of this effect may determine the pharmacodynamic features *in vivo*, i.e., therapeutic safety as well. Paradoxically, agents with low to moderate, rather than high, affinity may be desirable. Such findings triggered a reconsideration of the concept of uncompetitive antagonism of NMDA receptors in drug development [19, 22]. At present, many such agents are at different stages of development, e.g., carvedilol, ADCI, ES 242S, remacemide, felbamate, and budipine. On the other hand, uncompetitive NMDA receptor antagonists, such as amantadine and memantine - which fulfil the above criteria - have been used clinically for several years in the treatment of Parkinson's disease and dementia respectively, and do indeed rarely produce side effects at the therapeutic doses used in their respective indications.

In view of the above mentioned evidence, we have developed a series of novel uncompetitive NMDA receptor antagonists based on the 1-aminoalkylcyclohexane structure. The present study was devoted to compare the NMDA receptor antagonistic properties of these 1-aminoalkylcyclohexane derivatives in receptor-binding assays, patch clamp experiments, excitotoxicity *in vitro*, three convulsion models, and two models of motor impairment. The substi-

tutions of these 1-aminoalkylcyclohexanes are detailed in Table 6.

#### THE PRESENT INVENTION

5 It has now been found that certain 1-aminoalkylcyclohexanes have pronounced and unpredictable NMDA receptor antagonistic activity. Owing to the aforementioned property, the substances are suited for the treatment of a wide range of CNS disorders which involve disturbances of the glutamatergic transmission, preferably in the form of  
10 a pharmaceutical composition thereof wherein they are present together with one or more pharmaceutically-acceptable diluents, carriers, or excipients.


#### OBJECTS OF THE INVENTION

15 It is an object of the present invention to provide novel pharmaceutical compounds which are 1-aminoalkylcyclohexane NMDA receptor antagonists and pharmaceutical compositions thereof. It is a further object of the invention to provide a novel method of treating, eliminating, alleviating, palliating, or ameliorating undesirable  
20 CNS disorders which involve disturbances of glutamatergic transmission by the employment of such a compound of the invention or a pharmaceutical composition containing the same. An additional object of the invention is the provision of a process for producing the said 1-aminoalkylcyclohexane active principles. Yet additional objects will  
25 become apparent hereinafter, and still further objects will be apparent to one skilled in the art.

#### SUMMARY OF THE INVENTION

30 What we therefore believe to be comprised by our invention may be summarized inter alia in the following words:

A 1-aminoalkylcyclohexane compound selected from those of the formula



wherein  $R^*$  is  $-(CH_2)_n-(CR^6R^7)_m-NR^8R^9$

wherein  $n+m = 0, 1, \text{ or } 2$

wherein  $R^1$  through  $R^9$  are independently selected from hydrogen and lower-alkyl (1-6C), at least  $R^1$ ,  $R^4$ , and  $R^5$  being lower-alkyl;

5

such a compound wherein  $R^1$  through  $R^5$  are methyl;

such a compound wherein  $R^1$  is ethyl;

such a compound wherein  $R^2$  is ethyl;

such a compound wherein  $R^3$  is ethyl;

10

such a compound wherein  $R^4$  is ethyl;

such a compound wherein  $R^5$  is ethyl;

such a compound wherein  $R^5$  is propyl;

such a compound wherein  $R^6$  or  $R^7$  is methyl;

such a compound wherein  $R^6$  or  $R^7$  is ethyl; and

15

such a compound wherein the compound is selected from the group consisting of

1-amino-1,3,3,5,5-pentamethylcyclohexane,

1-amino-1,3,5,5-tetramethyl-3-ethylcyclohexane,

1-amino-1,5,5-trimethyl-3,3-diethylcyclohexane,


20

1-amino-1,5,5-trimethyl-cis-3-ethylcyclohexane,

1-amino-1,5,5-trimethyl-trans-3-ethylcyclohexane,

1-amino-1-ethyl-3,3,5,5-tetramethylcyclohexane,  
 1-amino-1-propyl-3,3,5,5-tetramethylcyclohexane,  
 N-methyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and  
 N-ethyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and  
 5 pharmaceutically-acceptable salts of any of the foregoing.

Moreover, a method-of-treating a living animal for  
 alleviation of a condition which is alleviated by an NMDA  
 receptor antagonist comprising the step of administering to  
 the said living animal an amount of a 1-aminoalkylcyclo-  
 10 hexane compound selected from those of the formula



wherein  $R^*$  is  $-(CH_2)_n-(CR^6R^7)_m-NR^8R^9$

wherein  $n+m = 0, 1, \text{ or } 2$

wherein  $R^1$  through  $R^9$  are independently selected from  
 hydrogen and lower-alkyl (1-6C), which is effective for  
 15 alleviation of the said condition;

such a method wherein  $R^1$  through  $R^5$  are methyl;  
 such a method wherein  $R^1$  is ethyl;  
 such a method wherein  $R^2$  is ethyl;  
 such a method wherein  $R^3$  is ethyl;  
 20 such a method wherein  $R^4$  is ethyl;  
 such a method wherein  $R^5$  is ethyl;

such a method wherein R<sup>5</sup> is propyl;

such a method wherein R<sup>6</sup> or R<sup>7</sup> is methyl;

such a method wherein R<sup>6</sup> or R<sup>7</sup> is ethyl; and

5 such a method wherein the compound is selected from the group consisting of

1-amino-1,3,3,5,5-pentamethylcyclohexane,

1-amino-1,3,5,5-tetramethyl-3-ethylcyclohexane,

1-amino-1,5,5-trimethyl-3,3-diethylcyclohexane,

1-amino-1,5,5-trimethyl-cis-3-ethylcyclohexane,

10 1-amino-1,5,5-trimethyl-trans-3-ethylcyclohexane,

1-amino-1-ethyl-3,3,5,5-tetramethylcyclohexane,

1-amino-1-propyl-3,3,5,5-tetramethylcyclohexane,


N-methyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and

N-ethyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and

15 pharmaceutically-acceptable salts of any of the foregoing;  
and

20 such a method wherein the compound is administered in the form of a pharmaceutical composition thereof comprising the compound in combination with one or more pharmaceutically-acceptable diluents, excipients, or carriers.

Further, an NMDA-receptor antagonist pharmaceutical composition comprising an effective NMDA-receptor antagonistic amount of a 1-aminoalkylcyclohexane compound selected from those of the formula





wherein R\* is  $-(CH_2)_n-(CR^6R^7)_m-NR^8R^9$

wherein  $n+m = 0, 1, \text{ or } 2$

5 wherein R<sup>1</sup> through R<sup>9</sup> are independently selected from hydrogen and lower-alkyl (1-6C), at least R<sup>1</sup>, R<sup>4</sup>, and R<sup>5</sup> being lower-alkyl, in combination with one or more pharmaceutically-acceptable diluents, excipients, or carriers;

such a pharmaceutical composition wherein R<sup>1</sup> through R<sup>5</sup> are methyl;

10 such a pharmaceutical composition wherein R<sup>1</sup> is ethyl;

such a pharmaceutical composition wherein R<sup>2</sup> is ethyl;

such a pharmaceutical composition wherein R<sup>3</sup> is ethyl;

such a pharmaceutical composition wherein R<sup>4</sup> is ethyl;

such a pharmaceutical composition wherein R<sup>5</sup> is ethyl;

15 such a pharmaceutical composition wherein R<sup>5</sup> is propyl;

such a pharmaceutical composition wherein R<sup>6</sup> or R<sup>7</sup> is methyl;

such a pharmaceutical composition wherein R<sup>6</sup> or R<sup>7</sup> is ethyl;

20 such a pharmaceutical composition wherein the compound is selected from the group consisting of

1-amino-1,3,3,5,5-pentamethylcyclohexane,

1-amino-1,3,5,5-tetramethyl-3-ethylcyclohexane,

1-amino-1,5,5-trimethyl-3,3-diethylcyclohexane,

25 1-amino-1,5,5-trimethyl-cis-3-ethylcyclohexane,

1-amino-1,5,5-trimethyl-trans-3-ethylcyclohexane,

1-amino-1-ethyl-3,3,5,5-tetramethylcyclohexane,


1-amino-1-propyl-3,3,5,5-tetramethylcyclohexane,

N-methyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and

N-ethyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and pharmaceutically-acceptable salts of any of the foregoing.


DETAILED DESCRIPTION OF THE INVENTION

5       The following details and detailed Examples are given by way of illustration only, and are not to be construed as limiting.



## Methods

## Chemistry



Preparation of 3-propyl-5,5-dimethyl-2-cyclohexene-1-one (1-7)

5 A solution of 3-ethoxy-5,5-dimethyl-2-cyclohexene-1-one [1] (5.04 g, 30 mmol) in ether was added dropwise to a stirred solution of propylmagnesium iodide prepared from 90 mg of magnesium and 90 mmol of 1-iodopropane in 60 ml of ether. After being stirred for 1h at ambient temperature, the reaction mixture was treated with 5% H<sub>2</sub>SO<sub>4</sub> solution. The organic phase was separated, washed with saline, dried  
10 over MgSO<sub>4</sub> and evaporated to give a crude oil which was separated on a silica gel column, eluting with hexane-ethyl acetate mixture. Cyclohexenone (1-7) was obtained as a colourless oil (2.0 g, 70%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.92 (3H, t, J=7 Hz); 1.03 (6H, s); 1.3 - 1.75 (2H, m); 2.16 (2H, t, J=7 Hz); 2.17 (2H, d, J=1.5 Hz); 2.21 (2H, s) and 5.87  
15 ppm (1H, t, J=1.5 Hz).

Such known cyclohexenones 1 were used to prepare compounds 2:

- 1-1 (R<sup>1</sup>=R<sup>2</sup>=R<sup>3</sup>=H) [commerc. available],
- 20 1-2 (R<sup>3</sup>=Me)\* [commerc. available],
- 1-3 (R<sup>2</sup>=R<sup>3</sup>=Me) [commerc. available],
- 1-4 (R<sup>1</sup>=R<sup>2</sup>=Me) [2],
- 1-5 (R<sup>1</sup>=R<sup>2</sup>=R<sup>3</sup>=Me) [commerc. available],
- 1-6 (R<sup>1</sup>=R<sup>2</sup>=Me, R<sup>3</sup>=Et) [3].
- 25 \*R<sup>n</sup>=H, if omitted

Other starting materials 1 are prepared in the same or similar manner.

General procedure for preparation of cyclohexanones 2.

30 Anhydrous copper (I) chloride (7.5 mmol) was added to a cooled solution of alkylmagnesium iodide (15-18 mmol) in ether. The mixture was stirred in an inert atmosphere for 5 minutes and a solution of 2-cyclohexene-1-one 1 (10 mmol)

in ether was added dropwise keeping the temperature below -5°C. After the addition of ketone was completed, the reaction mixture was stirred for 1 hour and carefully neutralized with saturated aqueous  $\text{NH}_4\text{Cl}$  solution. Traditional workup for Grignard reactions gave crude material which was separated on a silica gel column, eluting with a petroleum ether - ethyl acetate mixture. The cyclohexanones 2 were obtained as oils.

Yields and  $^1\text{H}$  NMR spectral data of compounds 2 are given in Table 1.

Such known cyclohexanones 2 were used to prepare compounds 3.

- 2-1 ( $\text{R}^4=\text{Me}$ )\* [commerc. available],  
2-2 ( $\text{R}^4=\text{Et}$ ) [4],  
2-3 ( $\text{R}^4=\text{Pr}$ ) [5],  
2-4 ( $\text{R}^3=\text{R}^4=\text{Me}$ ) [6],  
2-5 ( $\text{R}^3=\text{Me}$ ,  $\text{R}^4=\text{Et}$ ) [7],  
2-6 ( $\text{R}^3=\text{Me}$ ,  $\text{R}^4=\text{Pr}$ ) [8],  
2-7 ( $\text{R}^1=\text{R}^4=\text{Me}$ ) [9],  
2-8 ( $\text{R}^2=\text{R}^3=\text{R}^4=\text{Me}$ ) [10],  
2-9 ( $\text{R}^2=\text{R}^3=\text{Me}$ ,  $\text{R}^4=\text{Et}$ ) [11],  
2-13 ( $\text{R}^1=\text{R}^2=\text{R}^3=\text{R}^4=\text{Me}$ ) [commerc. available],  
2-14 ( $\text{R}^1=\text{R}^2=\text{R}^3=\text{Me}$ ,  $\text{R}^4=\text{Et}$ ) [10],  
2-15 ( $\text{R}^1=\text{R}^2=\text{R}^3=\text{Me}$ ,  $\text{R}^4=\text{Pr}$ ) [10].  
\* $\text{R}^n=\text{H}$ , if omitted.

Other intermediate cyclohexanones 2 are prepared in the same or a similar manner. Cyclohexanones 2 were used to prepare compounds 3:

General procedure for preparation of alkylcyclohexanols 3.

5 An ethereal solution of alkylmagnesium iodide (3-4 equivalents) was added dropwise to a cooled solution of cyclohexanone 2 in ether. The mixture was stirred for 1 hour at ambient temperature and carefully destroyed with saturated aqueous ammonium chloride. Traditional workup for Grignard reactions gave mixtures of diastereomeric alcohols 3, which were separated on a silica gel column eluting with petroleum ether - ethyl acetate.

10 Yields and <sup>1</sup>H NMR spectral data of compounds 3 are given in Table 2.

Such known cyclohexanols 3 were used to prepare compounds 4:

- 15 3-1 ((R<sup>3</sup>)(R<sup>4</sup>)=R<sup>5</sup>=Me)\* [9], i.e., R<sup>3</sup> or R<sup>4</sup> and R<sup>5</sup> are Me.  
3-4 (R<sup>3</sup>=R<sup>4</sup>=Me, R<sup>5</sup>=Me) [12],  
3-5 (R<sup>3</sup>=R<sup>5</sup>=Me, R<sup>4</sup>=Et) [13],  
3-7 (R<sup>1</sup>=R<sup>4</sup>=R<sup>5</sup>=Me) [14],  
3-8 (R<sup>1</sup>=R<sup>3</sup>=R<sup>4</sup>=R<sup>5</sup>=Me) [10],  
3-13 (R<sup>1</sup>=R<sup>2</sup>=R<sup>3</sup>=R<sup>4</sup>=R<sup>5</sup>=Me) [10],  
20 3-14 (R<sup>1</sup>=R<sup>2</sup>=R<sup>3</sup>=R<sup>4</sup>=Me, R<sup>5</sup>=Et) [15],  
\*R<sup>n</sup>=H, if omitted.

Other intermediate cyclohexenols 3 are prepared in the same or a similar manner.

25 General procedure for preparation of 1-alkyl-1-azidocyclohexanes 4.

The alcohol 3 was mixed with 1.7 - 2 N hydrazoic acid (10-13 equivalents) solution in chloroform, and cooled in an ice bath. A solution of TiCl<sub>4</sub> (1.2 equivalents) in chloroform was added dropwise while temperature was maintained below 5°C. The mixture was stirred at room  
30 temperature for 24 hours and passed down a column of

alumina, eluting with chloroform. Evaporation of solvent provided diastereomeric azides 4 which were purified by flash chromatography on silica gel, eluting with light petroleum ether.

5           Yields and <sup>1</sup>H NMR spectral data of compounds 4 are given in Table 3.

Other intermediate 1-alkyl-1-azidocyclohexanes 4 are prepared in the same or a similar manner.

10           Preparation of 1-nitromethyl-3,3,5,5-tetramethylcyclohexene (6).

15           A solution of 3,3,5,5-tetramethylcyclohexanone (2-13) (1.54 g, 10 mmol) and ethylenediamine (60 mg) in nitromethane (45 ml) was refluxed in argon atmosphere for 25 h. Excess of nitromethane was then removed in vacuo and the residue was purified by flash chromatography on silica gel, eluting with hexane - ethyl acetate (6:1). 1.2 g (61%) of 6 was obtained as an oil.

20           <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ 0.96 and 1.03 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.34 (2H, s, 4-CH<sub>2</sub>); 1.82 (2H, br s, 6-CH<sub>2</sub>); 4.80 (2H, s, CH<sub>2</sub>NO<sub>2</sub>) and 5.64 ppm (1H, br s, C=C-H).

25           Preparation of ethyl 3,3,5,5-tetramethylcyclohexylideneacetate (7).

30           To a stirred solution of triethyl phosphonoacetate (49.32 g, 0.22 mol) in dry THF (180 ml) under argon NaH (8.8 g, 0.22 mol, 60% suspension in mineral oil) was added in small portions while cooling with ice water. Stirring was continued for 1h at room temperature, then a solution of 3,3,5,5-tetramethylcyclohexanone (2-13) (30.85 g, 0.2 mol) was added over 10 min and the resulting mixture was refluxed for 22 h. It was then poured onto ice (400 g), the product was extracted with ether (4\*150 ml) and the solution dried over MgSO<sub>4</sub>. After concentration in vacuo an

oily residue was distilled at 145°C (11 mm) to give 36.8 g (86%) of 6 as an oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ 0.96 and 0.98 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.27 (3H, t, CH<sub>3</sub>-ethyl); 1.33 (2H, m, cyclohexane 4-CH<sub>2</sub>); 1.95 and 2.65 (total 4H, both s, cyclohexane 2,6-CH<sub>2</sub>); 4.14 (2H, q, CH<sub>2</sub>-ethyl) and 5.69 ppm (1H, s, =C-H).

Preparation of ethyl 3,3,5,5-tetramethylcyclohexylacetate (8).

Ethyl 3,3,5,5-tetramethylcyclohexylideneacetate (7) (4.48 g, 20 mmol) in ethanol (100 ml) was hydrogenated over 10% Pd/C (0.22 g, 5 wt.%) at 10 atm for 18 h. Filtration through Celite<sup>™</sup> and evaporation afforded 4.28 g (95%) of 8 as an oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ 0.89 and 1.02 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.26 (3H, t, J=7Hz, CH<sub>3</sub>-ethyl); 0.6-1.55 (7H, m, ring protons); 2.13 (2H, m, 2-CH<sub>2</sub>); and 4.12 ppm (2H, q, J=7Hz, CH<sub>2</sub>-ethyl).

Preparation of 2-methyl-(3,3,5,5-tetramethylcyclohexyl)-propan-2-ol (9).

A solution of ethyl 3,3,5,5-tetramethylcyclohexylacetate (8) (2.26 g, 10 mmol) in ether (20 ml) was added dropwise to a 2 M methylmagnesium iodide solution in ether (20 ml) over 15 min, while cooling with ice water. The mixture was refluxed for 2 h, cooled and quenched with saturated aqueous NH<sub>4</sub>Cl. After traditional workup the product was purified on silica gel column, eluting with a mixture of hexane-ethyl acetate (20:1) to give 1.7 g (80%) of 9 as an oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ 0.86 and 1.00 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.23 (6H, s, α-CH<sub>3</sub>); 1.36 (2H, d, J=5Hz, -CH<sub>2</sub>-); 0.6-2.04 ppm (8H, m, ring protons and OH).



Preparation of 2-methyl-(3,3,5,5-tetramethylcyclohexyl)-propyl-2-azide (10).

5 Boron trifluoride etherate (0.77 g, 0.69 ml, 5.44 mmol) was added dropwise to a stirred solution of 2-methyl-(3,3,5,5-tetramethylcyclohexyl)-propan-2-ol (9) (0.96 g, 4.53 mmol) and trimethylsilyl azide (0.63 g, 0.72 ml, 5.44 mmol) in benzene (10 ml). After being stirred for 24h at room temperature the mixture was poured into water (20 ml). The organic phase was separated and washed with saturated aqueous NaHCO<sub>3</sub> (10 ml) and saline (10 ml). The solution was dried over MgSO<sub>4</sub>, filtered and concentrated. The crude product was purified on silica gel column, eluting with hexane to give 0.56 g (52%) of 10 as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.87 and 1.01 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.27 (6H, s, α-CH<sub>3</sub>); 1.36 (2H, d, J=5Hz, -CH<sub>2</sub>-); 0.6-1.85 ppm (7H, m, ring protons).

Preparation of 2-(3,3,5,5-tetramethylcyclohexyl)-ethanol (11).

20 A solution of ethyl 3,3,5,5-tetramethylcyclohexyl-acetate 8 (1.8 g, 8.0 mmol) in ether (30 ml) was added dropwise to a stirred suspension of lithium aluminum hydride (0.9 g, 24.0 mmol) in ether (30 ml), which was cooled in an ice bath. The reaction mixture was refluxed for 3 h, cooled and residual lithium aluminum hydride was destroyed with water. The aqueous layer was separated and twice extracted with ether. The combined ether phases were washed with saline, dried over MgSO<sub>4</sub>, filtered and evaporated. The crude product was purified by flash chromatography on silica gel, eluting with hexane - ethyl acetate mixture (4 : 1) to give 1.2 g (79%) of 11 as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.89 and 1.00 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.44 (2H, q, J=7Hz, 2-CH<sub>2</sub>); 0.55-1.95 (8H, m, ring protons and OH) and 3.70 ppm (2H, t, J=7Hz, CH<sub>2</sub>O).

Preparation of 2-(3,3,5,5-tetramethylcyclohexyl)-ethyl methanesulfonate (12).

5 A solution of methanesulfonyl chloride (1.03 g, 0.7 ml, 9.0 mmol) in dry benzene (20 ml) was added to a stirred solution of 2-(3,3,5,5-tetramethylcyclohexyl)-ethanol (11) (1.1 g, 6.0 mmol) and triethylamine (1.2 g, 1.7 ml, 12 mmol) in benzene (40 ml), while cooling in an ice bath. The reaction mixture was stirred at room temperature for 3h, then filtered through a short silica gel column, 10 eluting with benzene. Evaporation of solvent gave 1.48 g (94%) of 12 as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.88 and 0.98 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.62 (2H, q, J=7Hz, 2-CH<sub>2</sub>); 0.65-2.0 (7H, m, ring protons) 3.0 (3H, s, CH<sub>3</sub>-SO<sub>2</sub>) and 4.29 ppm (2H, t, J=7Hz, CH<sub>2</sub>O).

15 Preparation of 2-(3,3,5,5-tetramethylcyclohexyl)-ethylazide (13).

The mixture of sodium azide (2.27 g, 34.2 mmol), 2-(3,3,5,5-tetramethylcyclohexyl)-ethylmethanesulfonate-(12) (1.46 g, 5.57 mmol) and dimethyl sulfoxide (20 ml) was 20 stirred at room temperature for 48 h, diluted with water (50 ml) and extracted with ether (3\*30 ml). The organic phase was washed with saline (30 ml), dried over MgSO<sub>4</sub>, filtered and evaporated. The crude product was purified by flash chromatography on silica gel, eluting with hexane to 25 give 0.93 g (80%) of (13) as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.87 and 0.99 (total 12H, both s, cyclohexane 3,5-CH<sub>3</sub>); 1.47 (2H, q, J=7Hz, 2-CH<sub>2</sub>); 0.55-1.9 (7H, m, ring protons) and 3.31 ppm (2H, t, J=7Hz, CH<sub>2</sub>N<sub>3</sub>).

30 Preparation of N-formyl-1,3,3,5,5-pentamethylcyclohexanamine (14-1).

To a vigorously stirred solution of 1,3,3,5,5-pentamethylcyclohexanol (3-13) (2.7 g, 15.6 mmol) and trimethylsilyl cyanide (2.36 g, 23.8 mmol) in acetic acid (2.5 ml) under argon 98% sulfuric acid (4.66 g, 47.6 mmol)

was added, keeping temperature below -5°C. The mixture was stirred at room temperature for 22 h, then it was poured onto ice (100 g), neutralised with 50% NaOH solution to pH ~ 7 and extracted with ether (3\*30 ml). The combined ether phases were washed with saline (50 ml), then dried over MgSO<sub>4</sub> and evaporated. A slightly yellow crystalline residue was treated with small amount of acetonitrile and filtered off to give 2.5 g (80%) of 14-1 as a white crystals, m.p. 104-106°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.91 and 0.93 (total 6H, both s, 3,5-CH<sub>3eq</sub>); 1.08 (2H, m, 2,6-CH<sub>eq</sub>); 1.13 and 1.15 (total 6H, both s, 3,5-CH<sub>3ax</sub>); 1.25 (2H, m, 4-CH<sub>2</sub>); 1.32 and 1.38 (total 3H, both s, 1-CH<sub>3</sub>); 1.70 and 2.12 (total 2H, both d, 14.7 Hz, 2,6-CH<sub>ax</sub>); 5.30 and 5.60 (total 1H, both br s, NH); 8.05 and 8.30 ppm (total 1H, both d, 2.0 and 12.7 Hz, resp., HCO).

Preparation of N-acetyl-1,3,3,5,5-pentamethylcyclohexanamine (14-2).

To a vigorously stirred solution of 1,3,3,5,5-pentamethylcyclohexanol (3-13) (3.0 g, 17.65 mmol) in acetonitrile (20 ml) fuming HNO<sub>3</sub> (6 ml) was added dropwise, keeping temperature below 45°C. The resulting mixture was stirred at 45-50°C for 6 h, then it was cooled, poured into water (30 ml) and neutralised with aqueous NH<sub>3</sub>. Aqueous phase was extracted with ether (3\*30 ml). The combined ether phases were washed with saline (30 ml), then dried over MgSO<sub>4</sub>, filtered and evaporated. The crude product was crystallised from cold acetonitrile to give 2.23 g (60%) of 14-2 as a white crystals, m.p. 110°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.90 and 1.12 (total 12H, both s, 3,5-CH<sub>3</sub>); 1.33 (3H, s, 1-CH<sub>3</sub>); 1.88 (3H, s, CH<sub>3</sub>C=O); 0.75-2.25 (6H, m. ring protons) and 5.3 ppm (1H, br s, NH).

Preparation of N-methoxycarbonyl-N,1,3,3,5,5-hexamethylcyclohexanamine (15).

Methyl chloroformate (0.97 g, 0.8 ml, 10.3 mmol) was added in one portion to a suspension of N,1,3,3,5,5-hexamethylcyclohexanamine hydrochloride (5-20) (1.13g, 5.13 mmol) and Na<sub>2</sub>CO<sub>3</sub> (1.63g, 15.4 mmol) in THF (30 ml). The resulting mixture was stirred at room temperature for 6h, and then it was diluted with water (50 ml) and extracted with ether (3\*30ml). The combined organic phases were washed with 10% K<sub>2</sub>SO<sub>4</sub>, saline, dried over MgSO<sub>4</sub>, filtered and evaporated. The crude product was purified by flash chromatography, eluting with hexane - ethyl acetate mixture (6 : 1) to give 0.90 g (78%) of (15) as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.93 and 1.07 (total 12H, both s, 3,5-CH<sub>3</sub>); 1.23 (3H, s, 1-CH<sub>3</sub>); 1.0-1.4 (4H, m, 4-CH<sub>2</sub> and 2,6-CH<sub>2eq</sub>); 2.56 (2H, d, J=14 Hz, 2,6-CH<sub>2ax</sub>); 2.87 (3H, s, CH<sub>3</sub>N) and 3.64 ppm (3H, s, CH<sub>3</sub>O).

Preparation of ethyl (3,3,5,5-tetramethylcyclohexylidene)cyanoacetate (16).

The mixture of 3,3,5,5-tetramethylcyclohexanone (2-13) (2.64g, 17 mmol), ethyl cyanoacetate (1.93, 17 mmol), acetic acid (0.2 ml) and ammonium acetate (0.2 g) in benzene (6.4 ml) was refluxed with a Dean-Stark apparatus for 10 h. To this benzene (30 ml) and saline (30 ml) was added, organic layer separated, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. The crude product was purified by flash chromatography, eluting with hexane to give 2.0 g (50%) of (16) as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 1.01 (6H, s, 3,5-CH<sub>3eq</sub>); 1.05 (6H, s, 3,5-CH<sub>3ax</sub>); 1.34 (3H, t, J=7Hz, ethyl-CH<sub>3</sub>); 1.42 (2H, s, 4-CH<sub>2</sub>); 2.46 and 2.79 (total 4H, both s, 2,6-CH<sub>2</sub>); and 4.29 ppm (2H, q, J=7Hz, CH<sub>2</sub>O).

Preparation of ethyl (1,3,3,5,5-pentamethylcyclohexyl)-cyanoacetate (17).

Anhydrous copper (I) chloride (0.8 g, 8 mmol) was added to a cooled solution of alkylmagnesium iodide (prepared from magnesium (0.46 g, 19.2 mmol) and iodo-

methane (2.84 g, 20 mmol) in ether (12 ml). The mixture was stirred in an inert atmosphere for 5 min and a solution of ethyl (3,3,5,5-tetramethylcyclohexylidene)cyanoacetate (16) (2 g, 8 mmol) in ether (10 ml), was added dropwise keeping the temperature below -15° C. After the addition of ketone was completed, the reaction mixture was stirred for 3 h and carefully neutralised with saturated aqueous NH<sub>4</sub>Cl solution. Traditional workup for Grignard reactions gave crude material which was separated on a silica gel column, eluting with a petroleum ether - ethyl acetate mixture (20:1) to give 1.0 g (47%) of 17 as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.98 (9H, s, 3,5-CH<sub>3eq</sub> and 1-CH<sub>3</sub>); 1.06 (6H, s, 3,5-CH<sub>3ax</sub>); 1.31 (3H, t, J=7Hz, ethyl-CH<sub>3</sub>); 1.2 - 1.5 (6H, m, ring protons); 3.41 (1H, s, α-CH) and 4.25 ppm (2H, q, J=7Hz, CH<sub>2</sub>O).

Preparation of 1-cyanomethyl-1,3,3,5,5-pentamethylcyclohexane (18).

The mixture of ethyl (1,3,3,5,5-pentamethylcyclohexyl)cyanoacetate (17) (1g, 3.7 mmol), LiCl (0.05g) and water (0.15 ml) in DMSO (2.5 ml) was heated at 150-160° C for 4h. Solution was poured into water (70 ml) and extracted with ether (4\*20 ml). Ether was washed with saline (2\*50 ml), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. Crude product was purified on silica gel column, eluting with a petroleum ether-ethyl acetate mixture (20:1) to give 0.66 g (94%) of 18 as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, TMS) δ: 0.98 (9H, s, 3,5-CH<sub>3eq</sub> and 1-CH<sub>3</sub>); 1.02 (6H, s, 3,5-CH<sub>3ax</sub>); 1.21 (3H, s, ring protons); 1.31 (3H, s, ring protons) and 2.31 ppm (2H, s, CH<sub>2</sub>CN). IR (neat) ν<sub>CN</sub>=2242 cm<sup>-1</sup>.

General procedure for preparation of alkylcyclohexanamine hydrochlorides 5-1-5-25.

A solution of 4, 10 or 13 - 15, 18 in ether was added dropwise to a stirred suspension of lithium aluminum hydride (4 equivalents) in ether, which was cooled in an

ice bath. The reaction mixture was stirred at room temperature in the case of 4, 10, 13 or refluxed in the case of 14, 15, 18 till complete conversion of starting material (TLC control). Residual lithium aluminum hydride was destroyed with water, the aqueous layer separated and twice extracted with ether. The combined ether phases were washed with saline, dried over NaOH, filtered and evaporated. The amine obtained was treated with HCl without characterization. The amine hydrochloride was prepared either by passing of HCl gas through the amine solution in hexane or by addition of a 1 N HCl solution in ether to the amine solution. In both cases the solvent was removed after HCl addition, the residue treated with hexane or acetonitrile and the crystalline product filtered off to give 5-1 - 5-25 with excellent purity.

The physical properties and yields of compounds 5-1 - 5-25 are given in Table 4.

<sup>1</sup>H NMR spectral data of compounds 5-1 - 5-25 are given in Table 5.

Additional 1-aminoalkylcyclohexanes and their hydrochlorides are prepared in the same or a similar manner. The hydrochlorides can be converted to the free base or other acid addition salts as disclosed under "ACID ADDITION SALTS".

Preparation of 3,3,5,5-tetramethylcyclohexylmethylamine hydrochloride (5-26).

A solution of 1-nitromethyl-3,3,5,5-tetramethylcyclohexene (6) (1.1 g, 5.63 mmol) in a mixture of ethanol (140 ml) and chloroform (2.8 ml) was hydrogenated over 10% Pd/C (280 mg) at 5 atm for 20 h, filtered and evaporated. The crude product was treated with ether, filtered and washed with ether to give 0.57 g (50%) of amine 5-26.

The physical properties and yield of compound 5-26 are given in Table 4.

<sup>1</sup>H NMR spectral data of compound 5-26 are given in Table 5.

5       Amine 5-27 was prepared according to the known procedure [16].

Amine 5-28 [17] was prepared according to the general procedure from corresponding azide [18]. All physical properties were in good agreement with data described [17].

10

\* \* \* \* \*

The purity of all compounds prepared was checked by GC (MN-OV-1, 25m\*0.53m, d<sub>f</sub> = 1.0µm, 50-270°C (10°C/min)).

\* \* \* \* \*

#### ACID ADDITION SALTS

15       As acids suitable for the formation of acid addition salts according to conventional procedure, there may be mentioned from the mineral series the following acids:

hydrochloric, hydrobromic, methanesulfonic, isothionic, sulfuric, phosphoric, and sulfamic acids and, from the  
20       organic series: acetic, propionic, maleic, fumaric, tartaric, citric, oxalic, and benzoic acids, to name a few. Preferred acids are hydrochloric, citric, and maleic. Other pharmaceutically-acceptable acid addition salts may be prepared, if desired, and one acid addition salt may be  
25       converted into another by neutralizing one salt, for example, the hydrochloride, resulting in the free base, and then reacidifying with a different selected mineral or organic acid, to prepare another pharmaceutically-acceptable acid addition salt, as is conventional in the art.

30

\* \* \* \* \*

**Table 1. Cyclohexanones 2**

Compd.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Yield (%)	<sup>1</sup> H - NMR (CDCl <sub>3</sub> , TMS) δ ppm
2-10	Me		Me	Pr	81.5	0.86 (3H, t, 6Hz); 0.98 (3H, s); 1.01 (3H, d, 5Hz); 1.05 - 1.35 (4H, m); 1.55 - 2.05 (4H, m); 2.11 (2H, s); 2.34 (1H, m)
2-11	Me	Me		Et	54	0.88 (3H, s); 0.90 (3H, t, 7Hz); 1.06 (3H, s); 1.15 - 1.45 (2H, m); 2.13 (2H, s); 1.45 - 2.45 (5H, m)
2-12	Me	Me		Pr	74	0.87 (6H, m); 1.15 (3H, s); 1.15 - 1.45 (4H, m); 2.13 (2H, s); 1.45 - 2.45 (5H, m)
2-16	Me	Me	Et	Et	83.5	0.78 (6H, t, 7Hz); 1.04 (6H, s); 1.37 (2H, q, 7Hz); 1.52 (2H, s); 2.16 (4H, s)
2-17	Me	Me	Pr	Pr	79	0.87 (6H, m); 1.03 (6H, s); 1.25 (8H, m); 1.53 (2H, s); 2.16 (4H, s)



Table 2. 1-Alkylcyclohexanols 3

Compd.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	Yield (%)	<sup>1</sup> H - NMR (CDCl <sub>3</sub> , TMS) $\delta$ ppm
3-2a				Et	Me	93	0.84 (3H, t, 7Hz); 1.17 (3H, s); 1.0 - 1.85 (12H, m)
3-2b			Et		Me		0.87 (3H, t, 7Hz); 1.21 (3H, s); 1.0 - 1.85 (12H, m)
3-3a				Pr	Me	93	0.86 (3H, t, 7Hz); 1.18 (3H, s); 1.0 - 1.9 (14H, m)
3-3b			Pr		Me		0.86 (3H, t, 7Hz); 1.19 (3H, s); 1.0 - 1.85 (14H, m)
3-6a				Me	Me	88	0.83 (3H, s); 0.86 (3H, m); 1.19 (3H, s); 1.0 - 1.85 (13H, m)
3-6b			Pr	Me	Me		0.88 (3H, t, 6.5Hz); 1.04 (3H, s); 1.17 (3H, s); 0.95 - 1.95 (13H, m)
3-9	Me			Me	Me	94	0.80 (3H, s); 0.81 (3H, t, 7Hz); 0.86 (3H, d, 6.5 Hz); 1.17 (3H, s); 0.9 - 2.0 (10H, m)
3-10	Me			Me	Me	88	0.81 (6H, m); 0.88 (3H, d, 6.5Hz); 1.17 (3H, s); 0.9 - 2.0 (12H, m)
3-11	Me	Me		Et	Me	84	0.87 (6H, m); 1.08 (3H, s); 1.18 (3H, s); 0.95 - 1.95 (10H, m)
3-12	Me	Me		Pr	Me	88	0.88 (6H, m); 1.09 (3H, s); 1.18 (3H, s); 0.9 - 1.95 (12H, m)
3-15	Me	Me	Me	Me	Pr	85	0.89 (9H, m); 1.21 (6H, s); 0.95 - 1.7 (11H, m)
3-16	Me	Me	Me (Et)	Et (Me)	Me	89	0.81 (3H, t, 7Hz); 0.89, 1.17 and 1.21 (total 12H, all s); 0.9 - 1.35 (5H, m); 1.35 - 2.0 (4H, m)
3-17	Me	Me	Me (Pr)	Pr (Me)	Me	88	0.84 (3H, m); 0.88 and 1.19 (total 12H, both s); 0.9 - 1.35 (7H, m); 1.35 - 2.0 (4H, m)
3-18	Me	Me	Et	Et	Me	87	0.78 (6H, t, 7Hz); 0.89 (3H, s); 1.19 (6H, s); 0.95 - 1.3 (7H, m); 1.3 - 2.05 (4H, m)
3-19	Me	Me	Pr	Pr	Me	90	0.86 (6H, t, 6.5); 0.88 (3H, s); 1.18 (6H, s); 0.9 - 1.3 (11H, m); 1.3 - 2.05 (4H, m)

Table 3. 1-Alkyl-1-azidocyclohexanes 4

Comp ound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	Yield (%)	<sup>1</sup> H - NMR (CDCl <sub>3</sub> , TMS) δ ppm
4-1a				Me	Me	31	0.89 (3H, d, 6.5Hz); 1.31 (3H, s); 0.95 - 2.0 (9H, m)
4-1b					Me	6	0.92 (3H, d, 6.5Hz); 1.28 (3H, s); 1.0 - 2.0 (9H, m)
4-2a				Et	Me	26	0.88 (3H, t, 7Hz); 1.29 (3H, s); 0.95 - 2.0 (11H, m)
4-2b					Me	4	0.88 (3H, t, 6.5Hz); 1.27 (3H, s); 1.0 - 2.0 (11H, m)
4-3a				Pr	Me	24	0.88 (3H, t, 6.5Hz); 1.29 (3H, s); 1.0 - 2.0 (13H, m)
4-3b					Me	11	0.88 (3H, t, 6.5Hz); 1.27 (3H, s); 1.0 - 2.0 (13H, m)
4-4				Me	Me	65	0.90 (3H, s); 1.08 (3H, s); 1.27 (3H, s); 1.0 - 1.95 (8H, m)
4-5				Me (Et)	Me	60	0.82 and 1.04 (total 3H, s); 0.82 (3H, t, 7Hz); 1.28 and 1.29 (total 3H, s); 0.9 - 2.0 (10H, m)
4-6				Me (Pr)	Me	66	0.85 and 1.07 (total 3H, s); 0.87 and 0.90 (total 3H, t, 6.5Hz); 1.29 (3H, s); 1.0 - 1.95 (12H, m)
4-7	Me (H)	H (Me)		Me (H)	Me	31	0.87 (6H, d, 6Hz); 1.27 and 1.29 (total 3H, s); 0.95 - 2.15 (8H, m)
4-8a	Me			Me	Me	42	0.86 (3H, d, 6Hz); 0.89 (3H, s); 1.09 (3H, s); 1.27 (3H, s); 0.95 - 1.9 (7H, m)
4-8b		Me		Me	Me	12	0.92 (3H, d, 6Hz); 0.94 (3H, s); 0.97 (3H, s); 1.36 (3H, s); 0.95 - 2.0 (7H, m)
4-9a	Me			Me	Me	47	0.81 (6H, s and m); 0.86 (3H, d, 6Hz); 1.27 (3H, s); 0.95 - 1.95 (9H, m)
4-9b		Me		Me	Me	12	0.81 (3H, t, 7Hz); 0.87 (3H, s); 0.91 (3H, d, 6Hz); 1.34 (3H, s); 0.95 - (9H, m)
4-10a	Me			Me	Me	44	0.81 (3H, s); 0.84 (3H, d, 6Hz); 0.87 (3H, m); 1.27 (3H, s); 1.0 - 2.0 (11H, m)
4-10b		Me		Me	Me	9	0.88 (6H, s and m); 0.91 (3H, d, 6Hz); 1.34 (3H, s); 1.0 - 1.95 (11H, m)
4-11a	Me	Me		Me	Me	45	0.91 (3H, t, 7Hz); 0.92 (3H, s); 1.12 (3H, s); 1.31 (3H, s); 1.0 - 1.9 (9H, m)
4-11b	Me	Me		Me	Me	12	0.92 (3H, t, 7Hz); 0.97 and 0.99 (total 6H, s); 1.37 (3H, s); 1.0 - 1.9 (9H, m)
4-12a	Me	Me		Pr	Me	54	0.90 (6H, s and m); 1.10 (3H, s); 1.28 (3H, s); 0.95 - 1.9 (11H, m)
4-12b	Me	Me		Pr	Me	7	0.89 (3H, t, 7Hz); 0.95 (3H, s); 0.98 (3H, s); 1.37 (3H, s); 1.0 - 1.9 (11H, m)
4-13	Me	Me		Me	Me	67	0.89 (6H, s); 1.18 (6H, s); 1.29 (3H, s); 0.95 - 1.9 (8H, m)
4-14	Me	Me		Me	Me	39	0.89 (6H, s); 0.96 (3H, t, 7Hz); 1.19 (6H, s); 1.0 - 1.9 (8H, m)
4-15	Me	Me		Me	Pr	65	0.89 (6H, s); 0.93 (3H, m); 1.18 (6H, s); 1.0 - 1.8 (10H, m)
4-16	Me	Me		Me (Et)	Me	77	0.82 (3H, m); 0.89, 1.14 and 1.18 (total 9H, s); 1.26 and 1.29 (total 3H, s); 0.95 - 1.9 (8H, m)
4-17	Me	Me		Me (Pr)	Me	71	0.86; 0.88 (total 3H, t, 6.5Hz); 0.90, 1.17; 1.19 (total 9H, s); 1.28; 1.32 (total 3H); 0.95-1.9 (10H, m)
4-18	Me	Me		Et	Me	66	0.78 (6H, t, 7Hz); 0.90 (3H, s); 1.18 (3H, s); 1.31 (3H, s); 0.95 - 1.95 (10H, m)
4-19	Me	Me		Pr	Me	61	0.89 (9H, s and m); 1.17 (3H, s); 1.27 (3H, s); 0.95 - 1.95 (14H, m)

**Table 4. Amino-cyclohexane derivatives 5**



Mrz 2/	Comp	Formula	M.W.	Elemental analysis						m.p. (°C)	Yield (%)
				Calculated (%)			Found (%)				
				C	H	N	C	H	N		
625	5-1a	C <sub>8</sub> H <sub>17</sub> N <sup>+</sup> HCl	163.72	58.7	10.5	8.6	58.7	10.5	8.6	> 250	63
631	5-1b	C <sub>8</sub> H <sub>17</sub> N <sup>+</sup> HCl	163.72	58.7	10.5	8.6	58.7	10.5	8.6	200 - 202	48
629	5-2a	C <sub>9</sub> H <sub>19</sub> N <sup>+</sup> HCl	177.75	60.8	10.8	7.9	60.8	10.8	7.9	> 250	66
630	5-2b	C <sub>9</sub> H <sub>19</sub> N <sup>+</sup> HCl	177.75	60.8	10.8	7.9	60.8	10.8	7.9	179 - 181	43
627	5-3a	C <sub>10</sub> H <sub>21</sub> N <sup>+</sup> HCl	191.78	62.6	11.1	7.3	62.6	11.1	7.3	> 250	80
628	5-3b	C <sub>10</sub> H <sub>21</sub> N <sup>+</sup> HCl	191.78	62.6	11.1	7.3	62.6	11.1	7.3	181 - 182	81
621	5-4	C <sub>9</sub> H <sub>19</sub> N <sup>+</sup> HCl	177.75	60.8	10.8	7.9	60.8	10.8	7.9	230 - 231	73
620	5-5	C <sub>10</sub> H <sub>21</sub> N <sup>+</sup> HCl	191.78	62.6	11.1	7.3	62.6	11.1	7.3	168 - 170	71
617	5-6	C <sub>11</sub> H <sub>23</sub> N <sup>+</sup> HCl	205.81	64.2	11.3	6.8	64.2	11.3	6.8	106 - 108	68
616	5-7	C <sub>9</sub> H <sub>19</sub> N <sup>+</sup> HCl	177.75	60.8	10.8	7.9	60.8	10.8	7.9	280 - 282	50
607	5-8a	C <sub>10</sub> H <sub>21</sub> N <sup>+</sup> HCl	191.78	62.6	11.1	7.3	62.6	11.1	7.3	>240	74
608	5-8b	C <sub>10</sub> H <sub>21</sub> N <sup>+</sup> HCl	191.78	62.6	11.1	7.3	62.6	11.1	7.3	>240	57
622	5-9a	C <sub>11</sub> H <sub>23</sub> N <sup>+</sup> HCl	205.81	64.2	11.3	6.8	64.2	11.3	6.8	250 - 253	68
624	5-9b	C <sub>11</sub> H <sub>23</sub> N <sup>+</sup> HCl	205.81	64.2	11.3	6.8	64.2	11.3	6.8	228 - 231	60
618	5-10a	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl	219.84	65.6	11.9	6.4	65.6	11.5	6.4	167 - 168	57
619	5-10b	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl	219.84	65.6	11.9	6.4	65.6	11.5	6.4	237 - 238	36
633	5-11a	C <sub>11</sub> H <sub>23</sub> N <sup>+</sup> HCl	205.81	64.2	11.3	6.8	64.2	11.3	6.8	255 - 257	69
632	5-11b	C <sub>11</sub> H <sub>23</sub> N <sup>+</sup> HCl	205.81	64.2	11.3	6.8	64.2	11.3	6.8	216 - 218	44
635	5-12a	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl	219.84	65.6	11.9	6.4	65.6	11.5	6.4	218 - 221	83
634	5-12b	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl	219.84	65.6	11.9	6.4	65.6	11.5	6.4	200 - 203	44
579	5-13	C <sub>11</sub> H <sub>23</sub> N <sup>+</sup> HCl	205.81	64.2	11.3	6.8	64.2	11.3	6.8	235 - 237	82
600	5-14	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl·H <sub>2</sub> O	237.86	60.6	10.6	5.9	60.6	10.6	5.9	215 - 218	74
601	5-15	C <sub>13</sub> H <sub>27</sub> N <sup>+</sup> HCl	233.87	66.8	11.7	6.0	66.8	11.7	6.0	> 280	88
615	5-16	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl	219.84	65.6	11.9	6.4	65.6	11.5	6.4	162 - 163	65
614	5-17	C <sub>13</sub> H <sub>27</sub> N <sup>+</sup> HCl·0.5H <sub>2</sub> O	242.84	64.3	12.0	5.8	63.8	12.0	5.6	106 - 107	54
623	5-18	C <sub>13</sub> H <sub>27</sub> N <sup>+</sup> HCl·H <sub>2</sub> O	251.89	62.0	10.8	5.6	62.0	10.8	5.6	99 - 102	78
626	5-19	C <sub>15</sub> H <sub>31</sub> N <sup>+</sup> HCl	261.93	68.8	12.0	5.3	68.8	12.0	5.3	167 - 169	72
640	5-20	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl	219.84	65.6	11.9	6.4	65.6	11.7	6.3	249-251	86
639	5-21	C <sub>13</sub> H <sub>27</sub> N <sup>+</sup> HCl	233.82	66.8	12.1	6.0	66.6	12.3	5.9	257-259	82
642	5-22	C <sub>13</sub> H <sub>27</sub> N <sup>+</sup> HCl·H <sub>2</sub> O	251.82	62.0	12.0	5.6	62.0	12.0	5.5	>210	98
645	5-23	C <sub>14</sub> H <sub>29</sub> N <sup>+</sup> HCl	247.85	67.8	12.2	5.7	67.6	12.3	5.6	205-207	89
644	5-24	C <sub>12</sub> H <sub>25</sub> N <sup>+</sup> HCl	219.84	65.6	11.9	6.4	65.4	11.9	6.2	>250	83
662	5-25	C <sub>13</sub> H <sub>27</sub> N <sup>+</sup> HCl·0.5H <sub>2</sub> O	242.84	64.3	12.0	5.8	64.9	11.9	5.7	>250	64
580	5-26	C <sub>11</sub> H <sub>23</sub> N <sup>+</sup> HCl	205.81	64.2	11.3	6.8	64.1	11.4	6.9	>230	50
557	5-27	C <sub>10</sub> H <sub>21</sub> N <sup>+</sup> HCl	191.75	62.6	11.6	7.3	62.3	11.6	7.2	>250(dec.)	70
641	5-28	C <sub>7</sub> H <sub>15</sub> N <sup>+</sup> HCl	149.7	56.2	10.8	9.4	55.9	11.0	9.2	283-285	69

Table 5 Spectral data of Amino-cyclohexane derivatives 5

Compd.	<sup>1</sup> H-NMR (CDCl <sub>3</sub> , TMS) δ, ppm
5-1a	0.89 (3H, d, 6Hz); 0.9 - 1.4 (3H, m); 1.44 (3H, s); 1.5 - 2.3 (6H, m); 8.3 (3H, br s)
5-1b	0.90 (3H, d, 5Hz); 1.48 (3H, s); 1.0 - 2.3 (9H, m); 8.3 (3H, br s)
5-2a	0.87 (3H, t, 7Hz); 1.45 (3H, s); 1.0 - 2.3 (11H, m); 8.35 (3H, br s)
5-2b	0.87 (3H, t, 7Hz); 1.46 (3H, s); 1.0 - 2.2 (11H, m); 8.3 (3H, br s)
5-3a	0.86 (3H, t, 6.5Hz); 0.95 - 1.4 (7H, m); 1.45 (3H, s); 1.5 - 2.2 (6H, m); 8.3 (3H, br s)
5-3b	0.85 (3H, t, 7Hz); 1.47 (3H, s); 0.95 - 2.2 (13H, m); 8.3 (3H, br s)
5-4	0.96 (3H, s); 1.05 (3H, s); 1.50 (3H, s); 1.1-1.95 (8H, m); 8.25 (3H, br s)
5-5	0.82 (3H, t, 7Hz); 0.90 and 1.04 (total 3H, both s); 1.48 and 1.50 (total 3H, both s); 1.1 - 2.0 (10H, m); 8.25 (3H, br s)
5-6	0.88 (3H, m); 0.94 and 1.07 (total 3H, both s); 1.49 and 1.52 (total 3H, both s); 1.1 - 2.0 (12H, m); 8.25 (3H, br s)
5-7	0.90 (6H, d, 6Hz); 1.44 and 1.50 (total 3H, both s); 0.95 - 2.4 (8H, m); 8.25 (3H, br s)
5-8a	0.90 and 0.91 (total 6H, d, s); 1.23 (3H, s); 1.44 (3H, s); 0.95 - 2.3 (7H, m); 8.2 (3H, br s)
5-8b	0.92, 0.96 and 0.98 (total 9H, d, s, s); 1.56 (3H, s); 1.04 - 2.4 (7H, m); 8.25 (3H, br s)
5-9a	0.83 (s) and 0.87 (m, total 9H); 1.47 (3H, s); 1.0 - 2.2 (9H, m); 8.15 (3H, br s)
5-9b	0.7 - 1.0 (m) and 0.89 (s, total 9H); 1.55 (3H, s); 1.05 - 2.2 (9H, m); (3H, br s)
5-10a	0.7 - 0.95 (m) and 0.86 (s, total 9H); 1.51 (3H, s); 0.85 - 2.3 (11H, m); 8.2 (3H, br s)

Table 5 (continued)

5-10b	0.7 - 1.0 (m) and 0.90 (s, total 9H); 1.54 (3H, s); 1.05 - 2.1 (11H, m); 8.2 (3H, br s)
5-11a	0.8 - 1.0 (m) and 0.91 (s, total 6); 1.22 (3H, s); 1.44 (3H, s); 1.0 - 2.3 (9H, m); 8.2 (3H, br s)
5-11b	0.88 (m) and 0.98 (s, total 9H); 1.50 (3H, s); 1.0 - 2.15 (9H, m); 8.2 (3H, br s)
5-12a	0.91 (6H, m); 1.22 (3H, s); 1.45 (3H, s); 1.0 - 2.3 (11H, m); 8.2 (3H, br s)
5-12b	0.89 (m); 0.97 (s, total 9H); 1.54 (3H, s); 1.0 - 2.2 (11H, m); 8.2 (3H, br s)
5-13	1.02 and 1.07 (total 12H, s); 1.26 (2H, m); 1.62 (3H, s); 1.71 (4H, m)
5-14	1.03 and 1.07 (total 12H, s); 1.09 (3H, t, 7Hz); 1.29 (2H, s); 1.59 and 1.81 (total 4H, d, 14Hz); 1.96 (2H, q, 7Hz); 8.15 (3H, br s)
5-15	0.93 (3H, t, 7Hz); 1.01 and 1.04 (total 12H, s); 1.29 (2H, s); 1.35 - 2.0 (4H, m); 1.70 (4H, m); 8.2 (3H, br s)
5-16	0.83 (3H, m); 1.00, 1.02 and 1.07 (total 9H, s); 1.2 - 1.5 (4H, m); 1.59 and 1.63 (total 3H, both s); 1.70 (4H, m); 8.25 (3H, br s)
5-17	0.87 (3H, m); 1.0 - 1.1 (9H, m); 1.1 - 1.4 (6H, m); 1.40 (4H, m); 1.59 (3H, s); 1.6 - 1.8 (4H, m); 8.25 (3H, br s)
5-18	0.78 (6H, t, 7Hz); 1.04 (6H, s); 1.27 (2H, m); 1.40 (4H, m); 1.59 (3H, s); 1.6 - 1.8 (4H, m); 8.25 (3H, br s)
5-19	0.87 (6H, m); 1.04 (6H, s); 1.1 - 1.5 (10H, m); 1.60 (3H, s); 1.5 - 1.95 (4H, m); 8.2 (3H, br s)
5-20	1.00 and 1.11 (total 12H, s); 1.29 (2H, m); 1.57 (3H, s); 1.72 (4H, dd, 14Hz); 2.56 (3H, t, 6Hz); 9.2 ppm (2H, br s)
5-21	0.98 and 1.11 (total 12H, s); 1.29 (2H, m); 1.58 (3H, t, 7Hz); 1.61 (3H, s); 1.82 (4H, m); 3.0 (2H, m); 9.1 ppm (2H, br s)
5-22	1.03 and 1.12 (total 12H, s); 1.32 (2H, m); 1.45 (3H, s); 1.64 and 1.97 (total 4H, d, 14Hz); 2.69 (6H, d, 5Hz)
5-23	0.85 (6H, s); 1.02 (6H, s); 0.6-1.95 (7H, m); 1.48 (6H, s); 1.60 (2H, d, 5Hz); 8.35 (3H, br s)
5-24	0.87 (6H, s); 0.98 (6H, s); 0.6-1.85 (9H, m); 3.02 (2H, m); 8.30 (3H, br s)
5-25	0.96(6H, s); 1.02(6H, s); 1.07(3H, s); 1.18(6H, s); 1.73 (2H, m); 3.03(2H, m); 8.28 (3H, br s)
5-26	0.97 (12H, br s); 1.0-2.2 (7H, m); 2.80 (2H, m); 8.35 ppm (3H, br s)
5-27	0.97 (6H, s); 1.04 (6H, s); 1.12 (1H, d, 13.7 Hz); 1.2 - 1.4 (5H, m); 1.92 (2H, d, 12.3 Hz); 3.47 (1H, m); 8.30 (3H, br s)
5-28	1.47 (3H, s) 1.2 - 2.2 (10H, m); 8.3 (3H, br s)

**Table 6 Basic Structure of the amino- and aminoalkylcyclohexanes.**
 $n+m=0,1,2$ 
 $R^1-R^9 = \text{lower alkyl}$ 

Mrz 2/	Compd	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>
625	5-1a	H	H	H	Me	Me	NH <sub>2</sub>
631	5-1b	H	H	Me	H	Me	NH <sub>2</sub>
629	5-2a	H	H	H	Et	Me	NH <sub>2</sub>
630	5-2b	H	H	Et	H	Me	NH <sub>2</sub>
627	5-3a	H	H	H	Pr	Me	NH <sub>2</sub>
628	5-3b	H	H	Pr	H	Me	NH <sub>2</sub>
621	5-4	H	H	Me	Me	Me	NH <sub>2</sub>
620	5-5	H	H	Me (Et)	Et (Me)	Me	NH <sub>2</sub>
617	5-6	H	H	Me (Pr)	Pr (Me)	Me	NH <sub>2</sub>
616	5-7	Me (H)	H (Me)	H (Me)	Me (H)	Me	NH <sub>2</sub>
643	5-8a	Me	H	Me	Me	Me	NH <sub>2</sub>
608	5-8b	H	Me	Me	Me	Me	NH <sub>2</sub>
622	5-9a	Me	H	Et	Me	Me	NH <sub>2</sub>
624	5-9b	H	Me	Me	Et	Me	NH <sub>2</sub>
618	5-10a	Me	H	Pr	Me	Me	NH <sub>2</sub>
619	5-10b	H	Me	Me	Pr	Me	NH <sub>2</sub>
633	5-11a	Me	Me	H	Et	Me	NH <sub>2</sub>
632	5-11b	Me	Me	Et	H	Me	NH <sub>2</sub>
635	5-12a	Me	Me	H	Pr	Me	NH <sub>2</sub>
634	5-12b	Me	Me	Pr	H	Me	NH <sub>2</sub>
579	5-13	Me	Me	Me	Me	Me	NH <sub>2</sub>
600	5-14	Me	Me	Me	Me	Et	NH <sub>2</sub>
601	5-15	Me	Me	Me	Me	Pr	NH <sub>2</sub>
615	5-16	Me	Me	Me (Et)	Et (Me)	Me	NH <sub>2</sub>
614	5-17	Me	Me	Me (Pr)	Pr (Me)	Me	NH <sub>2</sub>
623	5-18	Me	Me	Et	Et	Me	NH <sub>2</sub>
626	5-19	Me	Me	Pr	Pr	Me	NH <sub>2</sub>
640	5-20	Me	Me	Me	Me	Me	NHMe
639	5-21	Me	Me	Me	Me	Me	NHEt
642	5-22	Me	Me	Me	Me	Me	N(Me) <sub>2</sub>
645	5-23	Me	Me	Me	Me	H	CH <sub>2</sub> CM <sub>2</sub> NH <sub>2</sub>
644	5-24	Me	Me	Me	Me	H	CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>
662	5-25	Me	Me	Me	Me	Me	CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>
580	5-26	Me	Me	Me	Me	H	CH <sub>2</sub> NH <sub>2</sub>
557	5-27	Me	Me	Me	Me	H	NH <sub>2</sub>
641	5-28	H	H	H	H	Me	NH <sub>2</sub>

### PHARMACEUTICAL COMPOSITIONS

5           The active ingredients of the invention, together with one or more conventional adjuvants, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages thereof, and in such form may be employed as solids, such as coated or uncoated tablets or filled capsules, or liquids, such as solutions, suspensions, emulsions, elixirs, or capsules filled with the same, all for oral use; in the form of suppositories or capsules for  
10       rectal administration or in the form of sterile injectable solutions for parenteral (including intravenous or subcutaneous) use. Such pharmaceutical compositions and unit dosage forms thereof may comprise conventional or new ingredients in conventional or special proportions, with or  
15       without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. Tablets containing  
20       twenty (20) to one hundred (100) milligrams of active ingredient or, more broadly, ten (10) to two hundred fifty (250) milligrams per tablet, are accordingly suitable representative unit dosage forms.

### METHOD OF TREATING

25           Due to their high degree of activity and their low toxicity, together presenting a most favorable therapeutic index, the active principles of the invention may be administered to a subject, e.g., a living animal (including a human) body, in need thereof, for the treatment, alleviation, or amelioration, palliation, or elimination of  
30       an indication or condition which is susceptible thereto, or representatively of an indication or condition set forth elsewhere in this application, preferably concurrently, simultaneously, or together with one or more pharmaceutically-acceptable excipients, carriers, or diluents,

especially and preferably in the form of a pharmaceutical composition thereof, whether by oral, rectal, or parental (including intravenous and subcutaneous) or in some cases even topical route, in an effective amount. Suitable dosage ranges are 1-1000 milligrams daily, preferably 10-500 milligrams daily, and especially 50-500 milligrams daily, depending as usual upon the exact mode of administration, form in which administered, the indication toward which the administration is directed, the subject involved and the body weight of the subject involved, and the preference and experience of the physician or veterinarian in charge.

#### EXAMPLES OF REPRESENTATIVE PHARMACEUTICAL COMPOSITIONS

With the aid of commonly used solvents, auxiliary agents and carriers, the reaction products can be processed into tablets, coated tablets, capsules, drip solutions, suppositories, injection and infusion preparations, and the like and can be therapeutically applied by the oral, rectal, parenteral, and additional routes. Representative pharmaceutical compositions follow.

(a) Tablets suitable for oral administration which contain the active ingredient may be prepared by conventional tableting techniques.

(b) For suppositories, any usual suppository base may be employed for incorporation therein by usual procedure of the active ingredient, such as a polyethyleneglycol which is a solid at normal room temperature but which melts at or about body temperature.

(c) For parental (including intravenous and subcutaneous) sterile solutions, the active ingredient together with conventional ingredients in usual amounts are employed, such as for example sodium chloride and double-distilled water q.s., according to conventional procedure, such as



filtration, aseptic filling into ampoules or IV-drip bottles, and autoclaving for sterility.

Other suitable pharmaceutical compositions will be immediately apparent to one skilled in the art.

5           The following examples are again given by way of illustration only and are not to be construed as limiting.

EXAMPLE 1

Tablet Formulation

10           A suitable formulation for a tablet containing 10 milligrams of active ingredient is as follows:

	Mg.
15           Active Ingredient	10
Lactose	63
Microcrystalline	
Cellulose	21
Talcum	4
Magnesium stearate	1
20           Colloidal silicon	
dioxide	1

EXAMPLE 2

Tablet Formulation

25           Another suitable formulation for a tablet containing 100 mg is as follows:

	Mg.
30           Active Ingredient	100
Potato starch	20
Polyvinylpyrrolidone	10
Film coated and colored.	

The film coating material consists of:

	Lactose	100
	Microcryst. Cellulose	80
	Gelatin	10
5	Polyvinylpyrrolidone, crosslinked	10
	Talcum	10
	Magnesium stearate	2
	Colloidal silicon dioxide	3
10	Color pigments	5

---

EXAMPLE 3

## Capsule Formulation

A suitable formulation for a capsule containing 50 milligrams of active ingredient is as follows:

15		Mg.
	Active Ingredient	50
	Corn starch	20
20	Dibasic calcium phosphate	50
	Talcum	2
	Colloidal silicon dioxide	2

---

filled in a gelatin capsule.

EXAMPLE 4

## Solution for injection

A suitable formulation for an injectable solution containing one percent of active ingredient is as follows:

30	Active Ingredient	mg	12
	Sodium chloride	mg	8
	Sterile water to make ml		1

---

EXAMPLE 5

## Liquid oral formulation

A suitable formulation for 1 liter of a liquid mixture containing 2 milligrams of active ingredient in one milliliter of the mixture is as follows:

5

	G.
Active Ingredient	2
10 Saccharose	250
Glucose	300
Sorbitol	150
Orange flavor	10
Sunset yellow.	
15 Purified water to make a total of 1000 ml.	

EXAMPLE 6

## Liquid oral formulation

Another suitable formulation for 1 liter of a liquid mixture containing 20 milligrams of active ingredient in one milliliter of the mixture is as follows:

20

	G.
Active Ingredient	20
Tragacanth	7
Glycerol	50
Saccharose	400
30 Methylparaben	0.5
Propylparaben	0.05
Black currant-flavor	10
Soluble Red color	0.02
35 Purified water to make a total of 1000 ml.	

EXAMPLE 7

## Liquid oral formulation

Another suitable formulation for 1 liter of a liquid mixture containing 2 milligrams of active ingredient in one milliliter of the mixture is as follows:

5

	G.
Active Ingredient	2
10 Saccharose	400
Bitter orange peel tincture	20
Sweet orange peel tincture	15
Purified water to make a total of 1000 ml.	

15

EXAMPLE 8Aerosol formulation

180 g aerosol solution contain:

20

	G.
Active Ingredient	10
Oleic acid	5
Ethanol	81
25 Purified Water	9
Tetrafluoroethane	75

15 ml of the solution are filled into aluminum aerosol cans, capped with a dosing valve, purged with 3.0 bar.

EXAMPLE 9TDS formulation

100 g solution contain:

5		G.
	Active Ingredient	10.0
	Ethanol	57.5
	Propyleneglycol	7.5
10	Dimethylsulfoxide	5.0
	Hydroxyethylcellulose	0.4
	Purified water	19.6

15 1.8 ml of the solution are placed on a fleece covered by an adhesive backing foil. The system is closed by a protective liner which will be removed before use.

EXAMPLE 10Nanoparticle formulation

10 g of polybutylcyanoacrylate nanoparticles contain:

20		G.
	Active Ingredient	1.0
	Poloxamer	0.1
25	Butylcyanoacrylate	8.75
	Mannitol	0.1
	Sodiumchloride	0.05

30 Polybutylcyanoacrylate nanoparticles are prepared by emulsion polymersiation in a water/0.1 N HCl/ethanol mixture as polymerisation medium. The nanoparticles in the suspension are finally lyophilized under vacuum.

### PHARMACOLOGY - SUMMARY

5 The active principles of the present invention, and pharmaceutical compositions thereof and method of treating therewith, are characterized by unique advantageous and unpredictable properties, rendering the "subject matter as a whole", as claimed herein, unobvious. The compounds and pharmaceutical compositions thereof have exhibited, in standard accepted reliable test procedures, the following valuable properties and characteristics:

10 They are systemically-active, uncompetitive NMDA receptor antagonists with rapid blocking/unblocking kinetics and strong voltage dependency and are accordingly of utility in the treatment, elimination, palliation, alleviation, and amelioration of responsive conditions, by application or administration to the living animal host for  
15 the treatment of a wide range of CNS disorders which involve disturbances of glutamatergic transmission.

### PHARMACOLOGY

#### *In vitro*

20 Receptor Binding Studies

Male Sprague-Dawley rats (200-250g) were decapitated and their brains were removed rapidly. The cortex was dissected and homogenized in 20 volumes of ice-cold 0.32 M sucrose using a glass-Teflon homogenizer. The homogenate  
25 was centrifuged at 1000xg for 10 minutes. The pellet was discarded and the supernatant centrifuged at 20,000xg for 20 minutes. The resulting pellet was re-suspended in 20 volumes of distilled water and centrifuged for 20 minutes at 8000xg. Then the supernatant and the buffy coat were  
30 centrifuged three times (48,000xg for 20 minutes) in the presence of 50 mM Tris-HCl, pH 8.0. All centrifugation steps were carried out at 4°C. After resuspension in 5 volumes of 50 mM Tris-HCl, pH 8.0 the membrane suspension

was frozen rapidly at  $-80^{\circ}\text{C}$ . On the day of assay the membranes were thawed and washed four times by resuspension in 50 mM Tris-HCl, pH 8.0 and centrifugation at 48,000xg for 20 minutes. The final pellet was suspended in assay buffer. The amount of protein in the final membrane preparation was determined according to the method of Lowry with some modifications. The final protein concentration used for our studies was between 250-500  $\mu\text{g}/\text{ml}$ .

Membranes were re-suspended and incubated in 50 mM Tris-HCl, pH 8.0. Incubations were started by adding [ $^3\text{H}$ ]-(+)-MK-801 (23.9 Ci/mmol, 5nM) to vials with glycine (10 $\mu\text{M}$ ), glutamate (10 $\mu\text{M}$ ), and 0.1-0.25 mg protein (total volume 0.5 ml) and various concentrations of the agents tested (10 concentrations in duplicates). The incubations were continued at room temperature for 120 minutes, equilibrium always being achieved under the conditions used. Non-specific binding was defined by the addition of unlabeled MK-801 (10 $\mu\text{M}$ ). Incubations were terminated using a Millipore filter system. The samples were rinsed three times with 2.5ml ice cold assay buffer over glass fiber filters obtained from Schleicher & Schuell under a constant vacuum. Filtration was performed as rapidly as possible. Following separation and rinse, the filters were placed into scintillation liquid (5 ml; Ultima Gold) and radioactivity retained on the filters was determined with a conventional liquid scintillation counter (Hewlett Packard, Liquid Scintillation Analyser).

#### Patch Clamp

Hippocampi were obtained from rat embryos (E20 to E21) and were then transferred to calcium and magnesium free Hank's buffered salt solution (Gibco) on ice. Cells were mechanically dissociated in 0.05% DNAase / 0.3% ovomucoid (Sigma) following an 8 minute pre-incubation with 0.66% trypsin / 0.1% DNAase (Sigma). The dissociated cells were

then centrifuged at 18xg for 10 minutes, re-suspended in minimum essential medium (Gibco) and plated at a density of 150,000 cells cm<sup>-2</sup> onto poly-L-lysine (Sigma)-precoated plastic petri dishes (Falcon). The cells were nourished with NaHCO<sub>3</sub>/HEPES-buffered minimum essential medium supplemented with 5% fetal calf serum and 5% horse serum (Gibco) and incubated at 37°C with 5%CO<sub>2</sub> at 95% humidity. The medium was exchanged completely following inhibition of further glial mitosis with cytosine-β-D-arabinofuranoside (20μM Sigma) after about 7 days in vitro. Thereafter the medium was exchanged partially twice weekly.

Patch clamp recordings were made from these neurones with polished glass electrodes (4-6 mΩ) in the whole cell mode at room temperature (20-22°C) with the aid of an EPC-7 amplifier (List). Test substances were applied by switching channels of a custom-made fast superfusion system with a common outflow (10-20 ms exchange times). The contents of the intracellular solution were as follows (mM): CsCl (120), TEACl (20), EGTA (10), MgCl<sub>2</sub>(1), CaCl<sub>2</sub>-(0.2), glucose (10), ATP(2), cAMP (0.25); pH was adjusted to 7.3 with CsOH or HCl. The extracellular solutions had the following basic composition (mM): NaCl (140), KCl (3), CaCl<sub>2</sub> (0.2), glucose (10), HEPES (10), sucrose (4.5), tetrodotoxin (TTX 3\*10<sup>-4</sup>). Glycine (1μM) was present in all solutions: a concentration sufficient to cause around 80-85% activation of glycine<sub>α</sub> receptors. Only results from stable cells were accepted for inclusion in the final analysis, i.e., following recovery of responses to NMDA by at least 75% of their depression by the antagonists tested.

### Excitotoxicity

Cortical neurones were obtained from cerebral cortices of 17/18 day old fetal rats (Wistar), in general following the dissociation procedure described by [23]. After short trypsinization and gentle trituration with fire-polished



Pasteur pipettes, the cell suspension was washed by centrifugation. Cells were suspended in serum-free Neurobasal medium with B27 supplement (GIBCO) before plating onto poly-L-lysine (Sigma; 0.2mg/ml, 20h, 4°C) and laminin (Sigma; 2µg/ml, 1h, 37°C)-coated 96-well plates (Falcon, Primaria) at a density of  $5 \times 10^4$  cells/well. Cortical neurones were maintained at 37°C in humidified 10% CO<sub>2</sub>/90% air. One day after plating, 5µM cytosine-β-D-arabino-furanoside (Sigma) was added to each well for inhibition of glial cell proliferation. The medium was changed first, after 4 days in vitro and then every 4 days by replacing 2/3 of the medium with astrocyte-conditioned medium. Cortical neurones between day 12 and 14 in culture were used for the experiments.

New-born rat astrocytes were isolated non-enzymatically according to the method of [24]. Briefly, both hemispheres were dissected from 2-day-old rats, passed through an 80µm gauze, and triturated with Pasteur pipettes. Cell suspension was made in Dulbecco's modified essential medium (DMEM, Gibco) supplemented with 10% fetal calf serum (FCS, Hyclone), 2mM glutamine (Gibco) and 50µg/ml gentamycin and transferred into untreated, plastic culture flasks (Corning; 75cm<sup>3</sup>). Two days after plating the flasks were shaken for 10 minutes on a rotary platform (150 U/min) to remove microglial cells. The cultures were grown to confluency within 14 days, and the culture medium was changed twice weekly. Thereafter, the glial monolayers were extensively washed with serum-free Neurobasal medium (Gibco) to remove the serum. Flasks were then shaken several times to remove oligodendrocytes and neurones. To obtain conditioned medium from primary astrocytes, the cultures were incubated with fresh Neurobasal medium supplemented with B27 and glutamine. Every 2-3 days the conditioned medium was collected and replaced by fresh medium up to 4 times.

Exposure to EAA was performed in serum-free Neurobasal medium containing 100  $\mu$ M glutamate and the drug to be tested. After 20h of incubation, the cytotoxic effect was morphologically examined under a phase contrast microscope and biochemically quantified by measuring cell viability with the MTT test (Promega). This colorimetric assay measures the reduction of a tetrazolium component (MTT) into an insoluble formazan product by the mitochondria of living cells. After incubation of the cortical neurones with the dye solution for approximately 1-4 hours, a solubilization solution was added to lyse the cells and solubilize the colored product (incubation overnight at 37°C, 10% CO<sub>2</sub>, 90% RH). These samples were then read using an Elisa plate reader (Thermomax, MWG Biotech) at a wavelength of 570 nm. The amount of color produced was directly proportional to the number of viable cells.

#### *In vivo*

#### Anticonvulsive activity

NMR female mice (18-28 g) housed 5 per cage were used for the maximal electroshock (MES) and motor impairment tests. All animals were kept with water and food ad libitum under a 12 hour light-dark cycle (light on at 6 a.m.) and at a controlled temperature (20 $\pm$ 0.5°C). All experiments were performed between 10 a.m. and 5 p.m. Tested agents were injected 30 min. i.p. before the induction of convulsions if not stated otherwise (see below). All compounds were dissolved in 0.9% saline.

The MES test was performed together with tests for myorelaxant action (traction reflex) and motor coordination (rotarod). For the traction reflex test mice were placed with their forepaws on a horizontal rod and were required to place all 4 paws on the wire within 10 seconds. To test ataxia (motor coordination) mice were placed on rotarod (5 rpm) and were required to remain on the rod for 1 minute.

Only mice not achieving the criteria in all three repetitions of each test were considered to exhibit myorelaxation or ataxia respectively. These tests were followed by MES (100 Hz, 0.5 second shock duration, 50 mA shock intensity, 0.9 ms impulse duration, Ugo Basile) applied through corneal electrodes. The presence of tonic convulsions was scored (tonic extension of hind paws with minimum angle to the body of 90°). The aim was to obtain ED<sub>50</sub>s for all parameters scored (anticonvulsive activity and motor side effects) with use of the Litchfield Wilcoxon test for quantal dose responses. Division of the ED<sub>50</sub> for side effects (ataxia or myorelaxation) by the ED<sub>50</sub> for antagonism of electroshock convulsions was used as a therapeutic index (TI).

#### 15 Statistical analysis

IC<sub>50</sub>s in patch clamp, excitotoxicity, and binding studies were calculated according to the four parameter logistic equation using the Grafit computer program (Erithacus Software, England). Ki value for binding studies were then determined according to Cheng and Prusoff. Binding values presented are means ± SEM of 3-5 determinations (each performed in duplicate).

4-7 doses of antagonists were tested in each of the in vivo tests (5-8 animals per dose) to allow calculation of graded ED<sub>50</sub>s according to probit analysis (Litchfield and Wilcoxon) with correction for 0% to 100% effects. ED<sub>50</sub>s are presented with 95% confidence limits (Cl). Pearson product moment correlation analysis (Sigma Stat, Jandel Scientific) was used to compare in vitro potencies and in vivo anticonvulsant activity.

## RESULTS

### Binding

5 All cyclohexanes displaced [ $^3\text{H}$ ]-(+)-MK-801 binding to rat cortical membranes with  $\text{IC}_{50}\text{s}$  of between 4 and 150  $\mu\text{M}$  whilst  $\text{K}_\text{i}$  values as assessed with the Cheng-Prussoff equation were 2 fold lower (see Table 7).

### Patch Clamp

10 Steady-state inward current responses of cultured hippocampal neurones to NMDA (200 $\mu\text{M}$  with glycine 1 $\mu\text{M}$  at -70mV) were antagonized by the tested cyclohexanes with  $\text{IC}_{50}\text{s}$  of 1.3-99  $\mu\text{M}$  (Table 7). Peak and steady-state currents were affected to a similar degree making it unlikely that their effects were mediated at the glycine<sub>2</sub> site. Strong support for the uncompetitive nature of this antagonism was  
15 provided by the clear use- and voltage-dependency of their blockade. The weaker antagonists showed faster kinetics and stronger voltage-dependency.

### Excitotoxicity

20 Low  $\mu\text{M}$  concentrations of most cyclohexanes were effective neuroprotectants in vitro, with Mrz 2/579 seeming to be most potent in this regard (see Table 7). With most compounds full protection was obtained with 20 $\mu\text{M}$ .

### In vivo

#### Anticonvulsive activity

25 All cyclohexane derivatives inhibited MES-induced convulsions in mice with  $\text{ED}_{50}\text{s}$  ranging from 3.6 to 50 mg/kg i.p. (Table 7). Selected compounds were also tested against PTZ- and NMDA-induced convulsions (see [20,21] for methods) and showed comparable potency to the MES test  
30 (e.g., Mrz 2/579 had  $\text{ED}_{50}\text{s}$  in the PTZ- and NMDA tests of 5.5 and 3.7 mg/kg respectively). Their anticonvulsive potency was increased following i.v. administration (e.g., Mrz 2/579  $\text{ED}_{50}$  = 2.5 mg/kg). Mrz 2/579 was also active

5 following s.c. and somewhat less potent following p.o.  
administration ( $ED_{50}$ s of 4.6 and 13.7 mg/kg respectively).  
At doses within the anticonvulsive range, myorelaxation  
(traction test) and ataxia (rotarod test) were observed  
with some cyclohexanes. For the majority of them, no acute  
lethality was seen at up to 50mg/kg.

#### Correlation Analysis

10 There was a very good cross correlation between all  
three in vitro assays (all corr. coeffs.  $> 0.70$ ,  $p < 0.001$ ).  
There was also a good correlation between  
potencies in antagonizing NMDA-induced inward currents and  
protection against NMDA-induced toxicity in vitro with  
anticonvulsive activity in vivo (corr. coeffs.  $> 0.56$ ,  $p < 0.01$ ).




Table 7

Mrz 2/	[ <sup>3</sup> H] MK-801 IC <sub>50</sub> (μM) (μM)	SEM	Patch Clamp IC <sub>50</sub> (μM)	SEM	Glut. Tox. IC <sub>50</sub> (μM)	SD	MES ED <sub>50</sub> mg/kg	C.L.
557	17.6	0.9	18.5	2.7	6.7	2.0	43.9	35.6 - 54.1
579	1.9	0.1	1.3	0.02	2.2	0.0	3.6	2.2 - 6.1
580	15.9	0.8	12.9	0.4	5.6	0.8	27.3	12.8 - 64
600	2.4	0.1	3.7	0.2	2.1	0.2	22.6	43.0 - 197
601	7.4	0.7	10.5	0.8	3.5	0.3	15.6	10.4 - 23.4
607	8.2	0.3	13.8	1.5	10.1	2.2	22.9	18.3 - 28.7
608	6.6	1.3	12.7	1.2	16.6	1.1	20.6	18.3 - 23.2
614	13.6	1.3	13.9	1.9	> 10		23.5	15.7 - 34.9
615	2.5	0.1	2.9	0.1	2.3	0.1	6.1	3.4 - 10.7
616	15.0	0.4	34.2	4.6	9.1	2.1	24.0	15.6 - 36.8
617	51.8	3.9	57.4	7.3	> 70		54.9	42.9 - 70.4
618	32.7	2.4	43.7	9.4	17.6	2.9	24.0	9.6 - 59.5
619	72.1	6.7	60.8	5.4	30.9	2.9	44.6	32.0 - 62.3
620	32.2	2.1	99.0	10.4	38.4	1.6	41.3	32.9 - 51.7
621	36.7	4.4	92.4	19.0	> 100		36.9	22.6 - 60.3
622	15.0	0.6	64.8	11.7	19.3	8.1	21.0	16.1 - 27.5
623	3.3	0.2	3.7	0.7	4.5	0.6	13.1	9.9 - 17.2
624	15.4	1.2	31.0	3.6	2.7	0.6	47.2	41.8 - 53.2
625	46.8	8.1	244.9	40.1	39.4	6.3	129.8	42.5 - 395.6
626	11.6	1.5	9.6	2.0	19.0	3.3	41.2	29.9 - 56.7
627	70.3	3.3	209.7	1.0	26.6	5.7	43.9	30.3 - 63.7
628	35.6	4.4	125.5	0.8	27.3	4.5	73.2	33.6 - 159.4
629	39.4	2.4	218.6	1.6	> 300		58.5	38.3 - 89.2
630	44.3	3.8	> 100		> 100		> 30	
631	69.7	8.6	> 100		> 100		30.00	
632	2.0	0.2	6.4	0.6	10.9	0.4	11.04	7.7 - 15.8
633	6.8	0.5	13.9	3.2	5.4	0.9	8.78	3.6 - 21.4
634	15.5	1.0	10.8	2.6	19.0	3.5	> 30	
635	7.8	0.4	21.0	4.6	8.2	1.4	31.59	21.3 - 46.8
639	3.3	0.3	7.4	1.0	5.7	0.4	5.5	3.8 - 9.0
640	3.7	0.6	14.6	1.2	8.3	0.4	8.2	5.7 - 11.8
641	184.5	26.7	> 100		> 100		> 50	
642	10.2	1.6	42.5	6.5	29.3	3.3	8.04	5.1 - 12.7
643	3.6	0.5	13.5	1.7	12.0	0.9	18.65	10.8 - 32.2
644	3.8	3.7	4.1	1.8	4.3	0.4	52.98	27.8 - 100.8
645	85.1	30.6	20.4	3.6	> 100		65.61	43.8 - 98.2
Memantine	0.7	0.11	2.3	0.3	1.3	0.7	6.9	5.4 - 8.8
Amantadine	20.4	5.4	71.0	11.1	20.7	0.7	184.0	122 - 279
MK-801	0.0026	0.0002	0.14	0.10	0.012	0.002	0.16	0.13 - 0.21

Effects of cyclohexane derivatives and standard uncompetitive NMDA receptor antagonists on [<sup>3</sup>H]-(+)-MK-801 binding, NMDA induced currents in patch clamp experiments, glutamate toxicity in cultured cortical neurones and MES-convulsions in vivo. Binding K<sub>i</sub> values are means ±SEM of 3-5 experiments. IC<sub>50</sub>s (±SEM) in patch clamp and glutamate toxicity experiments were determined from data from at least 3 concentrations producing between 15 and 85% inhibition and at least 5 cells per concentration. For MES-induced convulsions, values are ED<sub>50</sub>s in mg/kg (95% confidence limits are shown in parentheses).

In addition, due at least in part to their amine substituent, the compounds of the present invention are also effective in non-NMDA indications, exhibiting immunomodulatory activity, antimalaria potency, anti-Borna virus activity, and anti-Hepatitis C activity.

\* \* \* \* \*

In conclusion, from the foregoing, it is apparent that the present invention provides novel, valuable, and unpredictable applications and uses of the compounds of the present invention, which compounds comprise the active principle according to the present invention, as well as novel pharmaceutical compositions thereof and methods of preparation thereof and of treating therewith, all possessed of the foregoing more specifically-enumerated characteristics and advantages.

The high order of activity of the active agent of the present invention and compositions thereof, as evidenced by the tests reported, is indicative of utility based on its valuable activity in human beings as well as in lower animals. Clinical evaluation in human beings has not been completed, however. It will be clearly understood that the distribution and marketing of any compound or composition falling within the scope of the present invention for use

in human beings will of course have to be predicated upon prior approval by governmental agencies, such as the U.S. Federal Food and Drug administration, which are responsible for and authorized to pass judgment on such questions.

5       **Conclusions**

          The presented 1-amino-alkylcyclohexanes represent a novel class of systemically-active, uncompetitive NMDA receptor antagonists with rapid blocking/unblocking kinetics and strong voltage-dependency. In view of their  
10       relatively low potency and associated rapid kinetics, they will be useful therapeutics in a wide range of CNS disorders which involve disturbances of glutamatergic transmission.

          These compounds accordingly find application in the  
15       treatment of the following disorders of a living animal body, especially a human. 1. Acute excitotoxicity such as ischaemia during stroke, trauma, hypoxia, hypoglycemia, and hepatic encephalopathy. 2. Chronic neurodegenerative diseases such as Alzheimer's disease, vascular dementia,  
20       Parkinson's disease, Huntington's disease, multiple sclerosis, amyotrophic lateral sclerosis, AIDS-neurodegeneration, olivopontocerebellar atrophy, Tourette's syndrome, motor neurone disease, mitochondrial dysfunction, Korsakoff syndrome, Creutzfeldt-Jakob disease. 3. Other  
25       disorders related to long term plastic changes in the central nervous system such as chronic pain, drug tolerance, dependence and addiction (e.g., opioids, cocaine, benzodiazepines, and alcohol). 4. Epilepsy, tardive dyskinesia, schizophrenia, anxiety, depression, acute pain,  
30       spasticity, and tinnitus. 5. In addition, as already stated, due at least in part to their amine substituent, the compounds of the present invention are also effective in non-NMDA indications, exhibiting immunomodulatory activity, antimalaria potency, anti-Borna virus activity,



and anti-Hepatitis C activity.

5

It is to be understood that the invention is not to be limited to the exact details of operation, or to the exact compositions, methods, procedures, or embodiments shown and described, as obvious modifications and equivalents will be apparent to one skilled in the art, and the invention is therefore to be limited only by the full scope which can be legally accorded to the appended claims.

### References


1. R.L. Frank, H.K. Hall (1950) J. Am. Chem. Soc. 72:1645-1648.
2. G.A. Hiegel, P. Burk. (1973) J. Org. Chem. 38:3637-3639.
3. N.F. Firrell, P.W. Hickmott. (1970) J. Chem. Soc. C:716-719.
4. G.H. Posner, L.L. Frye. (1984) Isr. J. Chem. 24:88-92.
5. G.L. Lemiere, T.A. van Osselaer, F.C. Anderweireldt. (1978) Bull. Soc. Chim. Belg. 87:771-782.
6. H.O. House, J.M. Wilkins. (1976) J. Org. Chem. 41:(25) 4031-4033.
7. A.R. Greenaway, W.B. Whalley. (1976) J. Chem. Soc. P.T. 1. :1385-1389.
8. S. Matsuzawa, Y. Horiguchi, E. Nakamura, I. Kuwajima. (1989) Tetrahedron 45:(2) 349-362.
9. H.O. House, W.F. Fischer. (1968) J. Org. Chem. 33:(3) 949-956.
10. Chiurdoglu, G., Maquestiau, A. (1954) Bull. Soc. Chim. Belg. 63: 357-378.
11. Zaidlewicz, M., Uzarewicz A., Zacharewicz, W. (1964) Roczniki Chem. 38: 591-597.
12. Crossley, A.W., Gilling, C. (1910) J. Chem. Soc. 2218.
13. Zaidlewicz, M., Uzarewicz, A. (1971) Roczniki Chem. 45: 1187-1194.
14. Lutz, E.T., van der Maas, J.H. (1982) Spectrochim. Acta, A. 38A: 283.
15. Lutz, E.T., van der Maas, J.H. (1981) Spectrochim. Acta, A. 37A: 129-134.

16. Ramalingam K., Balasubramanian, M., Baliah, V. (1972) Indian J. Chem. 10: 366-369.
17. Hamlin, K.E., Freifelder, M. (1953) J. Am. Chem. Soc. 75: 369-373.
- 5 18. Hassner, A., Fibinger, R., Andisik, D. (1984) J. Org. Chem. 49: 4237-4244.
19. W. Danysz, C.G. Parsons, I. Bresink, G. Quack (1995) Drug News Perspect. 8:261-277.
- 10 20. J.D. Leander, R.R. Lawson, P.L., Ornstein, D.M., Zimmerman (1988) Brain Res. 448:115-120.
21. C.G. Parsons, G. Quack, I. Bresink, L. Baran, E. Przegalinski, W. Kostowski, P. Krzascik, S. Hartmann, W. Danysz (1995). Neuropharmacology 34:1239-1258.
- 15 22. M.A. Rogawski (1993) Trends Pharmacol. Sci. 14:325-331.
23. Booher J. and Sensenbrenner M. (1972). Neurobiology 2:97-105.
24. Dichter, M. (1987) Brain Research 149:279.

We claim:

- 1 -

A 1-aminoalkylcyclohexane compound selected from those of the formula



5        wherein  $R^*$  is  $-(CH_2)_n-(CR^6R^7)_m-NR^8R^9$

wherein  $n+m = 0, 1, \text{ or } 2$

wherein  $R^1$  through  $R^9$  are independently selected from hydrogen and lower-alkyl (1-6C), at least  $R^1$ ,  $R^4$ , and  $R^5$  being lower-alkyl.

10

- 2 -

A compound of Claim 1 wherein  $R^1$  through  $R^5$  are methyl.

- 3 -

A compound of Claim 1 wherein  $R^1$  is ethyl.

- 4 -

15

A compound of Claim 1 wherein  $R^2$  is ethyl.

- 5 -

A compound of Claim 1 wherein  $R^3$  is ethyl.

- 6 -

A compound of Claim 1 wherein  $R^4$  is ethyl.

- 7 -

A compound of Claim 1 wherein R<sup>5</sup> is ethyl.

- 8 -

A compound of Claim 1 wherein R<sup>5</sup> is propyl.

5

- 9 -

A compound of Claim 1 wherein R<sup>6</sup> or R<sup>7</sup> is methyl.

- 10 -

A compound of Claim 1 wherein R<sup>6</sup> or R<sup>7</sup> is ethyl.

- 11 -

10

A compound of Claim 1 wherein the compound is selected from the group consisting of

1-amino-1,3,3,5,5-pentamethylcyclohexane,

1-amino-1,3,5,5-tetramethyl-3-ethylcyclohexane,

1-amino-1,5,5-trimethyl-3,3-diethylcyclohexane,

15

1-amino-1,5,5-trimethyl-cis-3-ethylcyclohexane,

1-amino-1,5,5-trimethyl-trans-3-ethylcyclohexane,

1-amino-1-ethyl-3,3,5,5-tetramethylcyclohexane,

1-amino-1-propyl-3,3,5,5-tetramethylcyclohexane,

N-methyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and

20

N-ethyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and


pharmaceutically-acceptable salts of any of the foregoing.

- 12 -

25

A method-of-treating a living animal for alleviation of a condition which is alleviated by an NMDA receptor antagonist, or for its immunomodulatory, antimalarial, anti-Borna virus, or anti-Hepatitis C effect, comprising the step of administering to the said living animal an amount of a 1-aminoalkylcyclohexane compound selected from those of the formula

- 51 -



wherein  $R^*$  is  $-(CH_2)_n-(CR^6R^7)_m-NR^8R^9$

wherein  $n+m = 0, 1, \text{ or } 2$

wherein  $R^1$  through  $R^9$  are independently selected from hydrogen and lower-alkyl (1-6C), which is effective for the said purpose.

5

- 13 -

A method of Claim 12 wherein  $R^1$  through  $R^5$  are methyl.

- 14 -

A method of Claim 12 wherein  $R^1$  is ethyl.

10

- 15 -

A method of Claim 12 wherein  $R^2$  is ethyl.

- 16 -

A method of Claim 12 wherein  $R^3$  is ethyl.

- 17 -

15

A method of Claim 12 wherein  $R^4$  is ethyl.

- 18 -

A method of Claim 12 wherein  $R^5$  is ethyl.

- 19 -

A method of Claim 12 wherein  $R^5$  is propyl.

20

- 20 -

A method of Claim 12 wherein  $R^6$  or  $R^7$  is methyl.

- 52 -

- 21 -

A method of Claim 12 wherein R<sup>6</sup> or R<sup>7</sup> is ethyl.

- 22 -

5 A method of Claim 12 wherein the compound is selected from the group consisting of


1-amino-1,3,3,5,5-pentamethylcyclohexane,  
1-amino-1,3,5,5-tetramethyl-3-ethylcyclohexane,  
1-amino-1,5,5-trimethyl-3,3-diethylcyclohexane,  
1-amino-1,5,5-trimethyl-cis-3-ethylcyclohexane,  
10 1-amino-1,5,5-trimethyl-trans-3-ethylcyclohexane,  
1-amino-1-ethyl-3,3,5,5-tetramethylcyclohexane,  
1-amino-1-propyl-3,3,5,5-tetramethylcyclohexane,  
N-methyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and  
N-ethyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and  
15 pharmaceutically-acceptable salts of any of the foregoing.

- 23 -

20 A method of Claim 12 wherein the compound is administered in the form of a pharmaceutical composition thereof comprising the compound in combination with one or more pharmaceutically-acceptable diluents, excipients, or carriers.

- 24 -

25 An NMDA-receptor antagonist pharmaceutical composition comprising an effective NMDA-receptor antagonistic amount, or an effective immunomodulatory, antimalarial, anti-Borna virus, or anti-Hepatitis C amount, of a 1-aminoalkylcyclohexane compound selected from those of the formula



wherein  $R^*$  is  $-(CH_2)_n-(CR^6R^7)_m-NR^8R^9$

wherein  $n+m = 0, 1, \text{ or } 2$

wherein  $R^1$  through  $R^9$  are independently selected from hydrogen and lower-alkyl (1-6C), at least  $R^1$ ,  $R^4$ , and  $R^5$  being lower-alkyl, in combination with one or more pharmaceutically-acceptable diluents, excipients, or carriers.

- 25 -

A pharmaceutical composition of Claim 24 wherein  $R^1$  through  $R^5$  are methyl.

- 26 -

A pharmaceutical composition of Claim 24 wherein  $R^1$  is ethyl.

- 27 -

A pharmaceutical composition of Claim 24 wherein  $R^2$  is ethyl.

- 28 -

A pharmaceutical composition of Claim 24 wherein  $R^3$  is ethyl.

- 29 -

A pharmaceutical composition of Claim 24 wherein  $R^4$  is ethyl.



- 30 -

A pharmaceutical composition of Claim 24 wherein R<sup>5</sup> is ethyl.

- 31 -

5 A pharmaceutical composition of Claim 24 wherein R<sup>5</sup> is propyl.

- 32 -

A pharmaceutical composition of Claim 24 wherein R<sup>6</sup> or R<sup>7</sup> is methyl.

10

- 33 -

A pharmaceutical composition of Claim 24 wherein R<sup>6</sup> or R<sup>7</sup> is ethyl.

- 34 -

15 A pharmaceutical composition of Claim 24 wherein the compound is selected from the group consisting of

1-amino-1,3,3,5,5-pentamethylcyclohexane,  
1-amino-1,3,5,5-tetramethyl-3-ethylcyclohexane,  
1-amino-1,5,5-trimethyl-3,3-diethylcyclohexane,  
1-amino-1,5,5-trimethyl-cis-3-ethylcyclohexane,  
20 1-amino-1,5,5-trimethyl-trans-3-ethylcyclohexane,  
1-amino-1-ethyl-3,3,5,5-tetramethylcyclohexane,  
1-amino-1-propyl-3,3,5,5-tetramethylcyclohexane,  
N-methyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and  
N-ethyl-1-amino-1,3,3,5,5-pentamethylcyclohexane, and  
25 pharmaceutically-acceptable salts of any of the foregoing.